

For high-risk NMIBC when
BCG does not work

STIL DOING IT MY WAY

With AD**STIL**ADRIN



Mike, an actual ADSTILADRIN patient.

Treatment Discussion Guide

Prepare for productive conversations with your doctor about high-risk NMIBC and ADSTILADRIN

INDICATION

ADSTILADRIN is a treatment for adults who have all the following:

- High-risk non-muscle-invasive bladder cancer (NMIBC)
- Cancer that is not responding to Bacillus Calmette-Guérin (BCG) therapy
- Carcinoma in situ (CIS) with or without other high-grade tumors

Please see Important Safety Information on page 4 and full **Prescribing Information**.

 **ADSTILADRIN[®]**
(nadofaragene firadenovec-vncg)
suspension, for intravesical use / 3×10^{11} viral particles/mL

Finding a treatment that fits your life starts with a conversation

If you have high-risk NMIBC or care for someone who does and BCG treatment isn't working, you may be wondering what's next.

Plan for a productive conversation with your doctor about NMIBC and treatment with ADSTILADRIN using the tips below, questions on page 3, and note sheet included at the end of this guide.

Treatment Discussion Tips



PREPARE BEFORE the appointment

Consider the following to help you plan your conversation:

- What type of treatment would best fit my lifestyle?
- How often will I need to come in for treatment?
- How might this treatment affect my current schedule and priorities?
- What types of side effects can I expect?
- Is financial assistance available?



ASK DURING the appointment

Write down your questions on the note sheet using the information above and on page 3.



REFLECT AFTER the appointment

- This treatment is my best way forward because...
- I will prepare for this treatment by...
- Questions to ask at my next appointment include...
- My next steps are...

Please see Important Safety Information on page 4 and full **Prescribing Information**.

 **ADSTILADRIN[®]**
(nadofaragene firadenovec-vncg)
suspension, for intravesical use / 3×10^{11} viral particles/mL

QUESTIONS to consider

Can you tell me more about my current diagnosis?

- How would you explain my current cancer stage?
- How has my cancer progressed?
- How is my current treatment working?

BCG isn't working, chemotherapy isn't right for me, and I want to keep my bladder.

- Do I still need to continue treatment with BCG?
- What are my next treatment options?

Is ADSTILADRIN right for me?

- Can ADSTILADRIN help me keep my bladder?
- How does ADSTILADRIN work?
- How well did ADSTILADRIN work for other people?

What can I expect during treatment with ADSTILADRIN?

- How should I prepare before each treatment session?
- How often will I receive treatment?
- What can I expect during treatment?
 - How will I be given ADSTILADRIN?
 - How long will my treatment sessions take?
 - Will I receive treatment in this same office?

How should I care for myself after each treatment?

- Will there be specific instructions or limitations for me?
- Can I resume my regular routine (work, family responsibilities, exercise, etc)?

When will I come back for follow-up appointments, and what can I expect during them?

- How will you check if treatment is working?
- Will I need any other procedures before my next dose?
- How long will I be prescribed ADSTILADRIN?

Please see Important Safety Information on page 4 and full [Prescribing Information](#).

 **ADSTILADRIN[®]**
(nadofaragene firadenovec-vncg)
suspension, for intravesical use / 3×10^{11} viral particles/mL

Learn more about ADSTILADRIN

Get helpful information
and resources, and
sign up for updates at
adstiladrin.com

Mike, an actual ADSTILADRIN patient.



Need help paying for treatment with ADSTILADRIN?

Call 1-844-NADOnow (1-844-623-6669)

Monday to Friday, 9 AM to 6 PM ET

or visit adstiladrin.com/financial-assistance/

IMPORTANT SAFETY INFORMATION

Who should not receive ADSTILADRIN?

Do not receive ADSTILADRIN if you have a sensitivity to interferon alfa or any of its components.

What is the most important information I should know about ADSTILADRIN?

Individuals who are immunosuppressed or immune-deficient should not prepare, administer, receive or come into contact with ADSTILADRIN.

What should I tell my healthcare provider?

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of ADSTILADRIN?

The most common side effects of ADSTILADRIN include:

Urinary discharge, fatigue, bladder spasm, urgency to urinate, and blood in your urine.

These are not all the possible side effects of ADSTILADRIN. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to FDA. Visit www.FDA.gov/medwatch or call 1-800-332-1088. You may also contact Ferring Pharmaceuticals at 1-888-FERRING.

What other information should I know about using ADSTILADRIN?

For 2 days following treatment, voided urine should be disinfected for 15 minutes with an equal volume of bleach before flushing.

For more important information, call 1-888-337-7464.

Please see full [Prescribing Information](#).



©2024 Ferring. ADSTILADRIN, Ferring and the Ferring Pharmaceuticals logo are trademarks of Ferring. US-ADST-2400170 v1 09/24

 **ADSTILADRIN[®]**
(nadofaragene firadenovec-vncg)
suspension, for intravesical use / 3×10^{11} viral particles/mL

